



ISSN: 2278-2648

International Journal of Research in Pharmacology & Pharmacotherapeutics (IJRPP)

IJRPP | Vol.12 | Issue 4 | Oct - Dec -2023

www.ijrpp.com

DOI : <https://doi.org/10.61096/ijrpp.v12.iss4.2023.254-263>

Research



Pharmacological screening and phytochemical evaluation of anti-diabetic activity of aloe vera in alloxan induced diabetic rats

Puppala. Sri Harshitha*, Dr. Sanapala Arun Kumar¹

Department Of Pharmacology, Sree Dattha Institute Of Pharmacy, Nagarjuna Sagar Road Sheriguda, Ibrahimpatnam Rangareddy - 501510.

*Author for Correspondence: Puppala. Sri Harshitha

Email: harshithasri76@gmail.com

	Abstract
Published on: 19 Oct 2023	<p>Diabetes mellitus is a most common endocrine disorder, affecting more than 300 million people worldwide. For these therapies developed along the principles of allopathic are often limited in efficacy, Carry the risk of adverse effects, and are often too costly, especially for the developing world. In order to identify complementary or alternative approaches to existing medications, we studied the anti-diabetic potential of leaves of <i>Aloe vera</i>. The acute oral toxicity studies of the extracts revealed no toxic effects up to the levels of 2000mg/kg b.wt. The aqueous and alcoholic extracts of 20 and 30mg/kg body weight of <i>Aloe vera</i> was screened for the presence of hypoglycemic and antidiabetic activity. In this study diabetes was induced by a single IP dose Alloxan monohydrate in 72hrs fasted rats. The FBGL was carried on 7th, 14th and 21st day and OGTT was measured on 8th, 15th and 22nd day. Glibenclamide was taken as the standard and the results are quite comparable with it. The studies were indicated that the leaves of <i>Aloe vera</i> are effective in regeneration of insulin secreting β-cells and thus possess antidiabetic activity. The aqueous and alcoholic extracts showed significant effect in decreasing the Fasting blood Glucose level and oral glucose tolerance test of rats and it's also showed good hypoglycemic activity in normal glycemic rats. The preliminary phytochemical analysis of the extracts of <i>Aloe vera</i> revealed the presence of Alkaloids, Tannins, Anthraquinones, Flavonoids, Saponins, Triterpenes, Sterols, Coumarin as the possible biologically active principles.</p>
Published by: DrSriram Publications	
<p>2023 All rights reserved.</p>  <p>Creative Commons Attribution 4.0 International License.</p>	<p>Keywords: <i>Aloe vera</i>, Alloxan monohydrate, Glibenclamide, FBGL and OGTT.</p>

INTRODUCTION

Diabetes Mellitus (DM)

Diabetes is one of the most common non-communicable diseases and a serious life-long condition appearing worldwide. The etiology of diabetes is a complex interaction of genetic and environmental factors. It is a heterogeneous group of metabolic disorders characterized physiologically by dysfunction of pancreatic beta cells and deficiency in insulin secretion or insulin activity and clinically by hyperglycemia or impaired glucose tolerance and other manifestable disorders. It is an endocrinological syndrome abnormally having high levels of sugar in the blood. This may be either due to insulin not being produced at all, is not made at sufficient levels, or is not as effective as it should be.

Diabetes is still a serious health problem all over the world since it is associated with increased morbidity and mortality rate. When compared with the general population, mortality and morbidity increase in diabetes is mainly due to the associated chronic complications both specific (microvascular) and nonspecific (macrovascular). Since the disease prevails in both genders and in all age groups, the general public has a concern about its control and treatment

Classification of DM

Diabetes is classified by underlying cause. The most common forms of diabetes are categorized as

Type 1, or insulin-dependent diabetes mellitus (IDDM) - an autoimmune disease in which the body's own immune system attacks the pancreatic beta cells, rendering it unable to produce insulin and

Type 2, or non-insulin-dependent diabetes mellitus (NIDDM) - in which there is resistance to the effects of insulin or a defect in insulin secretion.

Type 2 diabetes commonly occurs in adults associated with obesity. There are many underlying factors that contribute to the high blood glucose levels in these individuals. An important factor is the resistance to insulin in the body essentially ignoring its insulin secretions. A second factor is the decreased production of insulin by the cells of the pancreas. Therefore, an individual with Type 2 diabetes may have a combination of deficient secretion and deficient action of insulin. In contrast to Type 2 diabetes, Type 1 diabetes most commonly occurs in children and is a result of the body's immune system attacking and destroying the beta cells. The trigger for this autoimmune attack is not clear, but the result is the end of insulin production².

Multiple risk factors for the development of Type 2 diabetes mellitus:

- Family history (parents with diabetes).
- Obesity (i.e., $\geq 20\%$ over ideal body weight or body mass index $\geq 25\text{kg/m}^2$).
- Habitual physical inactivity.
- Impaired glucose tolerance.
- Hypertension ($\geq 140/90\text{mm Hg}$ in adults).
- High density lipoprotein (HDL) cholesterol $\leq 35\text{mg/dl}$ and/or triglyceride level $\geq 250\text{mg/dl}$.

History

The term "Diabetes" was first used around 250 B.C. It is a Greek word meaning "to syphon", reflecting how diabetes seemed to rapidly drain fluid from the affected individual. The Greek physician Aretaeus noted that affected individuals passed increasing amounts of urine as if there was "liquefaction of flesh and bones into urine". The complete term "diabetes mellitus" was coined in 1674 by Thomas Willis. Mellitus is Latin for honey, which is how Willis described the urine of diabetics.

Historical accounts reveal that as early as 700-200 BC, diabetes mellitus was a well recognized disease in India and was even distinguished as two types, a genetically based disorder and other one resulting from dietary indiscretion. Ancient Hindu writings document how black ants and flies were attracted to the urine of diabetics. The Indian physician Sushruta in 400 B.C. described the sweet taste of urine from affected individuals, and for many centuries to come, the sweet taste of urine was a key to the diagnosis.

Physicians have observed the effects of diabetes for thousands of years. One of the effects of diabetes is the presence of glucose in the urine (glucosuria). For much of the time, little was known about this fatal disease that caused weight loss of body, extreme thirst, and frequent urination. It was in 1922 that the first patient was successfully treated with insulin. Till the mid-1800s, the treatments offered for diabetes varied tremendously. A breakthrough in the puzzle of diabetes came in 1889. German physicians Joseph von Mering and Oskar Minkowski surgically removed the pancreas from dogs. The dogs immediately developed diabetes. Now that a link was established between the pancreas and diabetes, research focused on isolating the pancreatic extract that could treat diabetes. Dr. Frederick Banting succeeded in his experiments of isolating a pancreatic extract. The diabetic dog was kept alive for eight days by regular injections until supplies of the extract, at that time called "isletin", was exhausted. Experiments on dogs

showed that extracts from the pancreas caused a drop in blood sugar, caused glucose in the urine to disappear, and produced a marked improvement in clinical condition.

A young boy, Leonard Thompson, was the first patient to receive insulin treatment in the year 1922 and lived for thirteen years. Over the next 70 years, insulin was further refined and purified. A revolution came with the production of recombinant human DNA insulin in 1978. Instead of collecting insulin from animals, new human insulin could be synthesized. In 1923, Banting and Macloed were awarded the Nobel Prize for the discovery of insulin. In his Nobel Lecture, Banting concluded the following about their discovery: "Insulin is not a cure for diabetes; it is a treatment."

MATERIALS AND METHODS

The designing of methodology involves a series of steps taken in a systematic way in order to achieve the set goal(s) under the prescribed guidelines and recommendations. It includes in it all the steps from field trip to the observation including selection and collection of the medicinal plant, selection of dose value, standardization of protocol, usage of instruments, preparation of reagents, selection of specific solvents for extraction, formation of protocols and final execution of the standardized protocol. All this requires good build of mind and a good and soft technical hand to handle the materials and procedure in a true scientific manner.

Drugs and Chemicals

Drugs and Chemicals used in this study were of analytical grade and of highest purity procured from standard commercial sources in India.

Table 1: Drugs and Chemicals

<i>S.No</i>	<i>Materials</i>	<i>Company Name</i>
1.	Alloxan	Quali Kems Fine Chem Pvt, Ltd, Vadodara.
2.	Methanol	ChangshuYangyuan Chemicals, China.
3.	Alcohol	ChangshuYangyuan Chemicals, China.
4.	Glibenclamide	Orchid Pharma Ltd, Chennai.

Experimental animals

Healthy adult albino Wistar rats weighing 200-250grams of either sex were selected for the study. Animals were housed in appropriate cages in uniform hygienic conditions and fed with standard pellet diet (Amrul Laboratory Animal Diet) and water ad libitum. They were fasted overnight before the day of experiment, after 72hours of fasting from the day of Alloxan introduction. Animals were housed within the departmental animal house and the room temperature was maintained at 27° C. Animal studies had approval of IAEC.

Plant Material Collection

The leaves of *Aloe vera* were collected from the local market in Hyderabad in the month of January and was identified and authenticated from Department of Pharmacognosy. The plant material was cleaned, reduced to small fragments, air dried under shade at room temperature and coarsely powdered in a mixer. The powdered material was stored or taken up for extraction process.

Preparation of plant extracts

Preparation of Aqueous Extract

Dried leaves of *Aloe vera* were taken about 20gms into 250ml beaker containing 200ml of water. The contents were mixed well and then the mixture was boiled up to 80-90°C for 4-5hrs. Further the extract was filtered with whatmann filter paper. The filtrate was boiled until the concentrated residue is formed. The concentrated product was sealed in sample covers and stored under room temperature and used for further experiment to check the activities.

Preparation of Alcoholic Extract

Dried leaves of *Aloe vera* were taken about 20gms into 250ml beaker containing 200ml of Alcohol. The contents were mixed well and then the mixture was boiled up to 50-60°C for 4-5hrs. Further the extract was filtered with whatmann filter paper. The filtrate was boiled until the concentrated residue is formed. The concentrated product was sealed in sample covers and stored under room temperature and used for further experiment to check the activities.

Preliminary phytochemical analysis of the extracts

The extracts so obtained were subjected to preliminary phytochemical screening. Phytochemical studies were performed to identify the presence of various Phytoconstituents as follows:

Alkaloids

Extracts were dissolved individually in dilute Hydrochloric acid and filtered.

a. Mayer's Test: Filtrates were treated with Mayer's reagent (Potassium Mercuric Iodide). Formation of a yellow colored precipitate indicates the presence of alkaloids.

b. Wagner's Test: Filtrates were treated with Wagner's reagent (Iodine in Potassium Iodide). Formation of brown/reddish precipitate indicates the presence of alkaloids.

c. Dragendroff's Test: Filtrates were treated with Dragendroff's reagent (solution of Potassium Bismuth Iodide). Formation of red precipitate indicates the presence of alkaloids.

d. Hager's Test: Filtrates were treated with Hager's reagent (saturated picric acid solution). Presence of alkaloids confirmed by the formation of yellow colored precipitate.

Triterpenoids

a. Salkowski's Test: The extracts were treated with chloroform and filtered separately. The filtrate was treated with few drops of concentrated sulphuric acid, shaken and allowed to stand. If the lower layer turns red, sterols are present. If the lower layer turns golden yellow triterpenes are present.

Saponins

a. Froth Test: The extracts were diluted with distilled water to 20 ml and shaken in a graduated cylinder for 15 mins. The formation of 1 cm layer of foam indicates the presence of saponins.

b. Liberman Burchard Test: The extracts were treated with chloroform and filtered. The filtrates were treated with few drops of acetic anhydride boiled and cooled. Concentrated sulphuric acid was added through the sides of test tube. The formation of brown ring at the junction indicated the presence of steroidal saponins.

Flavonoids

a. Alkaline reagent Test: The extracts were treated with few drops of sodium hydroxide separately. Formation of intense yellow colour less on addition of few drops of dilute acid indicates the presence of Flavonoids.

b. Lead acetate Test: The extracts were treated with few drops of lead acetate solution. Formation of yellow precipitate indicates the presence of flavonoids.

Phenolic and Tannins

a. Ferric chloride Test: The extract was treated with few drops of neutral ferric chloride solution. The formation of bluish black colour indicates the presence of phenolics nucleus.

b. Gelatin Test: To the extract, 1% gelatin solution containing sodium chloride was added. The formation of white precipitate indicates the presence of tannins.

c. Vanillin hydrochloride Test: the extracts were treated with few drops of vanillin hydrochloride reagent. The conformation of pinkish red colour indicates the presence of tannins.

Selection of dose for animal study

The dose considered for the experiment on rats was obtained from conversion of human dose of *Aloe vera* (3-5 g/kg). The conversion factor of human dose (per 200 g body weight) is 0.018 for rats (Ghosh 1984). Hence the calculated dose for the rats (considering human dose 0.3 and 0.5 g/kg) is 20 and 30 mg/kg. Acute toxicity was done at dose of 2000mg/kg body weight.

Pharmacological evaluation

Preparation of extracts

The aqueous and alcoholic extracts of *Aloe vera* suspended in water in presence of 3%v/v Tween-80 solution. All the drugs were administered orally for experimental purpose. Each time preparations of the extracts were prepared when required. The drugs were administered at a constant volume of 10ml/kg for each animal.

Acute oral toxicity

The acute oral toxicity of aqueous and alcoholic extracts of *Aloe vera* was determined by using Albino Wistar rats (200-250g) which were maintained under standard conditions. The animals were fasted 12 hour prior to the experiment, up and down procedure OECD guideline no. 425 were adopted for toxicity studies. Animals were administered with single dose of individual extract up to 2000mg/kg and observed for its mortality during 2days and 7days study period (short term) toxicity and observed up to 7days for their mortality, behavioral and neurological profiles.

Assessment of Anti-diabetic Activity in Normal and Alloxan induced Rats

Assessment of hypoglycemic activity on normal rats.

Table 2: Group Classification

Group	Treatment	Dose(mg/kg)
Group 1	Normal control received distilled water	10ml/kg
Group 2	Standard group received Glibenclamide	10ml/kg
Group 3	Aqueous extract of <i>Aloe vera</i>	20 mg/kg
Group 4	Aqueous extract of <i>Aloe vera</i>	30 mg/kg
Group 5	Alcoholic extract of <i>Aloe vera</i>	20 mg/kg
Group 6	Alcoholic extract of <i>Aloe vera</i>	30 mg/kg

Procedure

Animals were divided randomly into six groups of four and each was fasted to overnight. The blood samples were withdrawn by tail vein at 0hour i.e. before I.P administration of extracts/standard/vehicle. Then blood was collected at an interval of 1, 2, 4, and 8 hour after the administration on 0th, 7th, 14th and 21st day respectively according to procedure blood glucose levels were measured by Glucometer (ONE TOUCH Glucometer).

Oral glucose tolerance test(OGTT) in normal rats

On the next day (1st, 8th, 15th and 22nd day) after the assessment of hypoglycemic activity OGTT was carried out in same normal animals.

Procedure

All the animals in each group were administered 2g/kg of glucose one hour after extract/ Glibenclamide/ vehicle administration. The blood samples were collected by tail vein at 0 hour, 0.5 hour, 1 hour, 1.5 hour and 2 hour after the administration of glucose load. Blood glucose levels were measured by Glucometer on 1st, 8th, 15th and 22nd day respectively.

Assessment of Anti-Diabetic Activity in Alloxan Induced Diabetic Rats

Induction of Diabetes

Albino Wistar rats of either sex weighing 200-250 g were selected for the study. All the animals were allowed free access to water and pellet diet and maintained at room temperature in rat cages.

Alloxan was dissolved in normal saline immediately before use. Diabetes was induced in 16 hour fasted rats by single intraperitoneal injection of 120 mg/kg body weight of freshly prepared alloxan in normal saline.

The rats after alloxanization were given 5% w/v glucose solution in feeding bottles for next 24 hours in their cages to prevent hypoglycemia. After 72 hours rats with fasting blood glucose levels greater than 200 mg/dl were selected and used for further studies

All the animals were observed for seven days for consistent hyperglycemia (fasting blood glucose level greater than 200 mg/dl and lesser than 400 mg/dl) and such animals were selected and divided into six groups of four each and used for the study of the following experimental models.

Table 3: Group Classification

Group	Treatment	Dose (mg/kg)
Group 1	Normal control received distilled water	10ml/kg
Group 2	Diabetic control received distilled water	10ml/kg

Group 3	Standard group received Glibenclamide	10ml/kg
Group 4	Aqueous extract of <i>Aloe vera</i>	20mg/kg
Group 5	Aqueous extract of <i>Aloe vera</i>	30mg/kg
Group 6	Alcoholic extract of <i>Aloe vera</i>	20 mg/kg
Group 7	Alcoholic extract of <i>Aloe vera</i>	30 mg/kg

Effect of Aqueous and Alcoholic extracts of *Aloe vera* on blood glucose levels in alloxan induced diabetic rats

All the animals of above groups were administered as per treatment protocol mentioned above. The blood samples were collected by retro orbital puncture at 0,1,2,4 and 8 hour after the administration. The treatment was continued for next 22 days. Again blood samples were also collected on 7th, 14th and 21st day after 1 hour administration for sub acute study. Blood glucose level was measured by Glucometer at various time intervals.

Oral glucose tolerance test (OGTT) in alloxan induced diabetic rats

On the 8th, 15th and 22nd day OGTT was carried out on the same alloxan induced diabetic animals used for assessment of anti-diabetic activity studies.

Procedure

All the animals in each group were administered 2g/kg of glucose one hour after extract/ Glibenclamide/ vehicle administration. The blood samples were collected by retro orbital puncture at 0 hour, 0.5 hour, 1 hour, 1.5 hour and 2 hour after the administration of the glucose load. The Blood samples were collected by tail vein and its blood glucose levels were measured by using a Glucometer apparatus.

Statistical analysis

The values were expressed as mean \pm SEM data was analyzed using one-way ANOVA followed by T-test. Two sets of comparison had made. i.e.

1. Normal control Vs All treated groups.
2. Diabetic Control Vs All treated groups.

Differences between groups were considered significant at $P < 0.001$ and $P < 0.05$ levels.

RESULTS AND DISCUSSION

Phytochemical screening of *Aloe vera*

The present investigation concluded that the isolated compounds from the plant *Aloe vera* shows the various Pharmacological effects was determined due to the presence of different phytochemical compound. Further study is needed for the isolation of the constituents present in the plant and its individual pharmacological activity should need to consider and ultimately it should be implemented for the benefit to human beings.

Table 4: Phytochemical screening of *Aloe vera*

S.No.	Phytoconstituents	Aqueous	Alcoholic
1.	Alkaloids	-	+
2.	Tannins	+	-
3.	Anthraquinones	+	+
4.	Flavonoids	-	+
5.	Saponins	-	-
6.	Triterpenes	+	-
7.	Sterols	+	+
8.	Coumarin	-	+

Acute toxicity testing

Acute toxicity studies revealed that the alcoholic extracts of *Aloe vera* were safe up to 2000 mg/kg of body weight and approximate LD 50 is more than 2000 mg/kg. No lethality or any toxic reactions was observed up to the end of the study period.

Hypoglycemic activity in normal rats

Fasting Blood Glucose Levels (FBGL) were within the range of 90-105 mg/dl in all the groups at 0 day. Repeated treatment with the doses of aqueous and alcoholic extract (100 and 200 mg/kg) significantly decrease the blood glucose level on 7th, 14th and 21st day, indicating that the extract produce significant hypoglycemic activity after repeated administration. Glibenclamide (10mg/kg) also significantly reduced Fasting Blood Glucose Level (FBGL) after repeated administration as compare to normal control group.

Repeated administration of both aqueous and alcoholic extracts had significantly ($p < 0.005$) reduced the FBGL on 7th, 15th and 21st day, indicating these extracts can produce hypoglycemia on repeated administration. However hypoglycemic activity was more significant on 7th, 14th and 21st day for Glibenclamide treated as compare with other groups. The results suggest that the both aqueous and alcoholic extracts possess significant hypoglycemic activity after repeated dose administration.

Effect of extracts of *Aloe vera* on fasting blood glucose level (FBGL) in normal rats

Table 5: Effect of extracts of *Aloe vera* on fasting blood glucose level (FBGL) in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level (mg/dl)		
		7 th day	14 th day	21 st day
Normal control	-	87.18±2.63	76.59±1.89	71.18±2.63
Glibenclamide	10	82.75±4.28	74.35±1.96	70.12±1.53
AQAV1	20	89.17±1.60	82.58±2.15	78.86±5.46
AQAV2	30	83.52±4.98	76.17±2.95	71.15±3.75
ALAV1	20	76.7±2.91	65.85±1.38	60.17±6.78
ALAV2	30	86.2±9.70	76.14±6.18	70.18±5.35

Values are expressed as mean± S.E.M. n=6. Significant values were compared with $p < 0.005$, normal control Vs all groups. Parent thesis indicates % reduction in BGL.

Oral glucose tolerance test (OGTT) -

Both the aqueous and alcoholic extracts of *Aloe vera* significantly ($P < 0.005$) suppress the rise in FBGL after glucose load (2g/kg) in rats, at first half-an-hour and up to 2hr time period as compare with other groups extract Glibenclamide on 8th, 15th and 22nd day. While aqueous and alcoholic extracts produced significant reduction in FBGL. Glibenclamide (10mg/kg) showed ($P < 0.005$) significant suppression in FBGL rise at first half-an-hour, 1hr and normalized FBGL within 2hr.

Table 6: Effect of extracts of *Aloe vera* on 8th, 15th and 22nd day in normal rats

Treatment	Dose (mg/kg)	Blood glucose level (mg/dl)		
		8 th day	15 th day	22 st day
Normal control	-	89.12±5.15	82.75±1.14	75.36±2.79
Glibenclamide	10	85.17±3.58	75.14±2.93	70.96±4.35
AQAV1	20	86.35±5.18	78.15±1.72	68.52±4.14
AQAV2	30	83.12±3.79	77.37±2.83	65.12±3.65
ALAV1	20	90.35±2.56	82.12±3.96	72.43±2.34
ALAV2	30	75.86±2.42	65.28±2.75	58.99±2.10

Values are expressed as mean ± S.E.M. n=6. Significant values were compared with $P < 0.005$. Normal control Vs all groups. Paranthesis indicates % reduction in BGL.

Anti-diabetic activity in alloxan induced diabetic rats

Changes in the fasting blood glucose levels in different groups are tabulated in Table No: 11. This data shown that blood glucose level of normal control animals has maintained throughout the study period.

The diabetic control group has shown significant increase in fasting blood glucose levels during this 21st day study period. Glibenclamide (10mg/kg) treated group has shown ($p < 0.05$) significant decrease in fasting blood glucose level during 7th, 14th and 21st day of study period.

Effect of *Aloe vera* extracts on antidiabetic activity in alloxan induced diabetic rats

The animals treated with 100 and 200mg/kg of aqueous and alcoholic of different extracts shown significant decrease ($P<0.05$) in FBGL on 7th, 14th and 21st day of treatment when compare to other groups of animals. The aqueous extracts have reduced more (%) in FBGL when compared to alcoholic extracts except standard group. The detailed results are summarized in Table No: 12

Table 7: Effect of extracts of *Aloe vera* on fasting blood glucose level (FBGL) in Alloxan induced diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level (mg/dl)		
		7 th day	14 th day	21 st day
Normal control	-	85.70±6.90	75.82±5.15	68.36±2.98
Diabetic control	10	296.15±21.53	272.52±12.15	265.58±24.39
Glibenclamide	10	263.10±25.19	246.14±85.28	228.14±52.76
AQAV1	20	375.29±68.17	360.35±15.85	332.74±12.63
AQAV2	30	381.26±15.89	368.24±19.85	351.23±21.96
ALAV1	20	292.15±86.75	271.15±59.13	221.85±36.15
ALAV2	30	223.52±16.85	185.15±02.61	153.75±55.89

Values are expressed as mean ± S.E.M. n=6. Significant values were compared with $P<0.05$. Normal control Vs all groups. Paranthesis indicates % reduction in BGL.

Oral glucose tolerance test (OGTT) on 8th, 15th and 22nd day-

Both the aqueous and alcoholic extracts of *Aloe vera* are significantly ($P<0.05$) suppress the rise in FBGL after glucose load (2g/kg) in rats, at first half-an-hour and up to 2hr time period as compare with other groups extract Glibenclamide on 8th, 15th and 22nd day. While aqueous and alcoholic extracts produced significant reduction in FBGL. Glibenclamide (10mg/kg) showed ($P<0.05$) significant suppression in FBGL rise at first half-an-hour, 1hr and normalized FBGL within 2hr.

Table 8: Effect of extracts of *Aloe vera* on 8th, 15th and 22nd day in Diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level (mg/dl)		
		8 th day	15 th day	22 st day
Normal control	-	85.12±2.85	75.28±1.91	63.14±2.89
Diabetic control	10	283.25±11.71	251.14±20.95	221.39±19.86
Glibenclamide	10	365.89±75.50	286.15±39.52	275.93±15.78
AQAV1	20	262.13±72.89	232.71±25.53	198.17±13.99
AQAV2	30	283.82±10.27	262.13±10.78	242.89±15.32
ALAV1	20	264.18±93.56	221.80±96.15	186.55±11.89
ALAV2	30	363.12±10.28	321.18±25.98	282.15±19.12

Values are expressed as mean ± S.E.M. n=6. Significant values were compared with $P<0.05$. Normal control Vs all groups. Paranthesis indicates % reduction in BGL.

CONCLUSION

The study was performed to find out the beneficial effects of two different extracts of leaves of *Aloe vera* in normoglycaemic rats and alloxan induced diabetic rats and the results reveal that the plant has beneficial effects on blood glucose levels.

In current scenario, herbs are the potent sources of medicines used in the treatment of various disease and disorders. Since, plants are used as medicine there is prompt need of evaluation of plant species, therefore, the present work was conceived to evaluate the phytochemical and pharmacological screening of leaves of *Oregano*. Alkaloids, Tannins, Anthraquinones, Flavonoids, Saponins, Triterpenes, Sterols and Coumarin.

The aqueous and alcoholic extracts had hypoglycemic activity because the presence of flavonoids which are rich in treatment of hypoglycaemia with less side effects. Flavonoids might be producing hypoglycaemic effect by a mechanism independent from insulin secretion e.g. by the inhibition of endogenous glucose production or by the inhibition of intestinal glucose absorption. The present study *Aloe vera* of both aqueous and alcoholic extracts was showed significant effect on glucose tolerance and also showed reduction in fasting blood glucose levels in normal diabetic rats.

The data of the blood glucose level of rats treated with Alloxan (150mg/kg body weight) produced diabetes within 72 hours. After 72 hours of Alloxan administered the blood glucose levels of rats were observed. It was observed that significant lowering of sugar in aqueous and alcoholic extract. The administration of different extracts at a dose of 20 and 30 mg/kg showed significant anti-hyperglycaemic effect at 22nd day which was evident from the 7th day onwards as compared to standard. The aqueous and alcoholic extract of *Aloe vera* has showed better anti-hyperglycaemic effect of the extract on the fasting blood sugar levels on diabetic rats are shown in table. The decreasing blood glucose levels are comparable with that of 10 mg/kg of Glibenclamide. The Glibenclamide (10 mg/kg body weight) shows significant effect on compare to the initial and more significant effect on the 22nd Day compare to the initial. The aqueous and alcoholic extracts of 20 and 30mg/kg body weight shows significant ($P < 0.05$), effect.

Results of anti-diabetic activity in normal and alloxan induced rats the extracts established the scientific basis for the utility of these plants in the treatment of diabetes. The extracts have shown significant reduction in blood glucose levels in normal and alloxan induced diabetic rats and produced maximum anti-diabetic activity and are higher than the hypoglycemic activity of Glibenclamide in the diabetic rats. In glucose loaded animals, the drug has reduced the blood glucose to the normal levels. It is possible that the drug may be acting by potentiating the pancreatic secretion or increasing the glucose uptake. In conclusion, these extract showed significant anti-diabetic effect in normal and diabetic rats after administration. Thus the claim made by the traditional Indian systems of medicine regarding the use of these plants in the treatment of diabetes stands confirms.

REFERENCES

1. Consultation, W. H. O. Definition, diagnosis and classification of diabetes mellitus and its complications. Vol. 1. Part, 1999.
2. Alberti, Kurt George Matthew Mayer, and PZ ft Zimmet. "Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation." *Diabetic medicine* 15.7 (1998): 539-553.
3. Gress, Todd W., et al. "Hypertension and antihypertensive therapy as risk factors for type 2 diabetes mellitus." *New England Journal of Medicine* 342.13 (2000): 905-912.
4. Ahmed, Awad M. "History of diabetes mellitus." *Saudi medical journal* 23.4 (2002): 373-378.
5. Zimmet, Paul Z., Daniel J. McCarty, and Maximilian P. de Courten. "The global epidemiology of non-insulin-dependent diabetes mellitus and the metabolic syndrome." *Journal of Diabetes and its Complications* 11.2 (1997): 60-68.
6. American Diabetes Association. "Type 2 diabetes in children and adolescents." *Pediatrics* 105.3 (2000): 671-680.
7. Saltiel, Alan R., and C. Ronald Kahn. "Insulin signalling and the regulation of glucose and lipid metabolism." *Nature* 414.6865 (2001): 799-806.
8. Browning, Jeffrey D., and Jay D. Horton. "Molecular mediators of hepatic steatosis and liver injury." *Journal of Clinical Investigation* 114.2 (2004): 147-152.
9. Aronoff, Stephen L., et al. "Glucose metabolism and regulation: beyond insulin and glucagon." *Diabetes Spectrum* 17.3 (2004): 183-190.
10. Ross, E. J., and D. C. Linch. "Cushing's syndrome—killing disease: discriminatory value of signs and symptoms aiding early diagnosis." *The Lancet* 320.8299 (1982): 646-649.
11. Levin, Marvin E., Vincenza C. Boisseau, and Louis V. Avioli. "Effects of diabetes mellitus on bone mass in juvenile and adult-onset diabetes." *New England Journal of Medicine* 294.5 (1976): 241-245.
12. Sherwin, Robert, and Philip Felig. "Pathophysiology of diabetes mellitus." *The Medical clinics of North America* 62.4 (1978): 695-711.
13. Faber, O. K., and C. Binder. "C-peptide response to glucagon: a test for the residual β -cell function in diabetes mellitus." *Diabetes* 26.7 (1977): 605-610.
14. Ioannidis, Ioannis. "Pathophysiology of Type 1 diabetes." *Diabetes in Clinical Practice: Questions and Answers from Case Studies* 31 (2007): 23.

15. Scheen, A. J. "Pathophysiology of type 2 diabetes." *Acta Clinica Belgica* 58.6 (2003): 335-341.
16. Vambergue, A., et al. "[Pathophysiology of gestational diabetes]." *Journal de gynecologie, obstetrique et biologie de la reproduction* 31.6 Suppl (2002): 4S3-4S10.
17. Inzucchi, Silvio E. "Diagnosis of diabetes." *New England Journal of Medicine* 367.6 (2012): 542-550.
18. Turner, R. C., et al. "Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33)." *lancet* 352.9131 (1998): 837-853.
19. Brownlee, M., and A. Cerami. "The biochemistry of the complications of diabetes mellitus." *Annual review of biochemistry* 50.1 (1981): 385-432.
20. Pari, L., and G. Saravanan. "Antidiabetic effect of Cogent db, a herbal drug in alloxan-induced diabetes mellitus." *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 131.1 (2002): 19-25.